

M. J. Drake's research while affiliated with The University of Arizona and other places

Overview

Publications (281)

Thermal infrared observations and thermophysical characterization of OSIRIS-REx target asteroid (101955) Bennu

Article

- May 2014
- [Joshua Emery](#)
- [Y.R. Fernández](#)
- [M.S.P. Kelley](#)
- [...]
- [Julie E. Ziffer](#)

Near-Earth Asteroids (NEAs) have garnered ever increasing attention over the past few years due to the insights they offer into Solar System formation and evolution, the potential hazard they pose, and their accessibility for both robotic and human spaceflight missions. Among the NEAs, carbonaceous asteroids hold particular interest because they ma...

Cite

[Request full-text](#)

Partition coefficients at high pressure and temperature

Article

- Jan 2014
- [Kevin Righter](#)
- [Lisa R. Danielson](#)
- [M.J. Drake](#)
- [K. Domanik](#)

Cite

[Request full-text](#)

Accretion disk origin of Earth's water

Article

- Jun 2013

- [Luca Vattuone](#)
- [Marco Smerieri](#)
- [Letizia Savio](#)
- [...]
- [Mario Rocca](#)

Earth's water is conventionally believed to be delivered by comets or wet asteroids after the Earth formed. However, their elemental and isotopic properties are inconsistent with those of the Earth. It was thus proposed that water was introduced by adsorption onto grains in the accretion disc prior to planetary growth, with bonding energies so high...

[Cite](#)

[Request full-text](#)

[A first-principles characterization of water adsorption on forsterite grains](#)

[Article](#)

- Jun 2013
- [Abu Md Asaduzzaman](#)
- [Slimane Laref](#)
- [P A Deymier](#)
- [...]
- [M J Drake](#)

Numerical simulations examining chemical interactions of water molecules with forsterite grains have demonstrated the efficacy of nebular gas adsorption as a viable mechanism for water delivery to the terrestrial planets. Nevertheless, a comprehensive picture detailing the water-adsorption mechanisms on forsterite is not yet available. Towards this...

[Cite](#)

[Request full-text](#)

[Thermal and Physical Characterization of the OSIRIS-REx Target Asteroid \(101955\) 1999 RQ36](#)

[Article](#)

- Oct 2012
- [Joshua Emery](#)
- [M. S. Kelley](#)
- [Y. R. Fernandez](#)
- [...]

- [M. J. Drake](#)

The OSIRIS-REx mission, the third in NASA's New Frontier line, will launch in 2016, visit the near-Earth asteroid (101955) 1999 RQ36, and return samples of its regolith to Earth in 2023. Ground-based observations have already revealed a great deal about 1999 RQ36, including the spectral type (B-type), size, and rotation period. To further character...

[Cite](#)

[Request full-text](#)

Electronic Structure Calculations of Water Adsorption on Forsterite Grains: Implications for Planetary Water

[Article](#)

- Sep 2011

- [K. Muralidharan](#)

- [M. J. Drake](#)

[Cite](#)

[Request full-text](#)

Accretion Disk Origin of Earth's Water: Laboratory Experiments

[Article](#)

- Sep 2011

- [Luca Vattuone](#)

- [Marco Smerieri](#)

- [Letizia Savio](#)

- [...]

- [Mario Rocca](#)

[Cite](#)

[Request full-text](#)

Experimental Evidence of Adsorption as a Source of Planetary Water

[Article](#)

- Mar 2011

- [Luca Vattuone](#)
- [Mario Rocca](#)
- [K. Muralidharan](#)
- [...]
- [M. J. Drake](#)

Many Earth oceans of water could have been adsorbed onto grains in the accretion disk prior to the formation of the terrestrial planets.

[Cite](#)

[Request full-text](#)

Computer simulations of water interactions with low-coordinated forsterite surface sites: Implications for the origin of water in the inner solar system

[Article](#)

- Nov 2010
- [Helen E King](#)
- [M. Stimpfl](#)
- [P Deymier](#)
- [...]
- [Nora H. de Leeuw](#)

Adsorption of water to fractal dust grains during accretion has been proposed as a possible source of water for rocky planets. We have used computer simulations to study the feasibility of chemisorption onto forsterite dust grains by investigating the adsorption of dissociated water to stoichiometric and defective surfaces. Defects were modeled usi...

[Cite](#)

[Request full-text](#)

Where on Earth has our water come from?

Article

Full-text available

- Oct 2010
- [Nora H. de Leeuw](#)
- [Richard Richard A Catlow](#)
- [Helen E King](#)
- [...]
- [Michael J Drake](#)

The presence of water in the Earth has long been an enigma. However, computer modelling techniques have shown that the adsorption of water onto the fractal surfaces of interplanetary dust particles, which are present in the planetary accretion disk, is sufficiently strong to provide a viable origin of terrestrial water.

Cite

[Download full-text](#)

Asteroid (101955) 1999 RQ36: Optimum Target for an Asteroid Sample Return Mission

Article

- Jul 2010
- [D. S. Lauretta](#)
- [M. J. Drake](#)
- [R. P. Benzel](#)
- [...]
- [D. J. Scheeres](#)

[Cite](#)

[Request full-text](#)

Atomistic Simulations of Adsorption of Water onto Forsterite and Fayalite Planar Surfaces: Implication for the Origin of Water in the Inner Solar System

Article

- Mar 2010
- M. Stimpfl
- K. Muralidharan
- Nora H. de Leeuw
- [...]
- M. J. Drake

Using atomistic and electronic structure calculations, it is shown that adsorption is a significant source of terrestrial planetary water.

Cite

[Request full-text](#)

Subsolidus Metal --- Olivine Trace Element Partitioning

Article

- Mar 2010
- Eddy Hill
- G. R. Huss
- K. Domanik
- M. J. Drake

Metal - olivine subsolidus diffusion experiments provide diffusion rates and trace-element diffusion profiles that allow us to study the cooling rates of mesosiderites and calculate subsolidus distribution coefficients .

Cite

[Request full-text](#)

Thermophysical Characterization of Potential Spacecraft Target (101955) 1999 RQ36

Article

- Mar 2010

- [Joshua Emery](#)
- [Y. R. Fernández](#)
- [M. S. Kelley](#)
- [...]
- [H. Campins](#)

We report on thermal emission measurements of 1999 RQ36 from Spitzer. The derived size is in agreement with radar measurements, and we find a moderately high thermal inertia and homogeneous surface properties.

[Cite](#)

[Request full-text](#)

Metal/Silicate Partitioning of the Moderately Siderophile Elements: The Effect of Temperature and C Concentration

[Article](#)

- Mar 2010
- [Eddy Hill](#)
- [K. Domanik](#)
- [M. J. Drake](#)

Increasing T in a magma ocean would result in reduced metal/silicate partitioning for Mo, W and V, simultaneously, the C-concentration in the metal may determine the presence or absence of V in the metal-melt.

[Cite](#)

[Request full-text](#)

Crystallization of magmatic iron meteorites: The role of mixing in the molten core

[Article](#)

- Feb 2010
- [Nancy Chabot](#)
- [Michael J. DRAKE](#)

Abstract— The IIIAB group is the largest of the magmatic iron meteorite groups and consequently is commonly used to test models of asteroid core crystallization. Simple fractional crystallization calculations appear to reproduce the general shape of the

elemental trends observed in the IIIAB group when these trends are plotted vs. Ni, as is traditi...

[Cite](#)

[Request full-text](#)

Crystallization of magmatic iron meteorites: The effects of phosphorus and liquid immiscibility

[Article](#)

- Feb 2010
- [Nancy Chabot](#)
- [Michael J. DRAKE](#)

Abstract— Magmatic iron meteorites are commonly thought to have formed by fractional crystallization of the metallic cores of asteroid-sized bodies. As fractional crystallization proceeds, light elements such as P and S become enriched in the molten portion of the core. The light element content of the metallic liquid influences the partitioning be...

[Cite](#)

[Request full-text](#)

A magma ocean on Vesta: Core formation and petrogenesis of eucrites and diogenites

[Article](#)

- Feb 2010
- [Kevin Righter](#)
- [MICHAEL J. DRAKE](#)

Abstract— Available evidence strongly suggests that the HED (howardite, eucrite, diogenite) meteorites are samples of asteroid 4 Vesta. Abundances of the moderately siderophile elements (Ni, Co, Mo, W and P) in the HED mantle indicate that the parent body may have been completely molten during its early history. During cooling of a chondritic compo...

[Cite](#)

[Request full-text](#)

An experimental study of Ag and Pd partitioning between solid and liquid metal, with applications to iron meteorites

[Article](#)

- Feb 2010
- [Nancy Chabot](#)

- [MICHAEL J. DRAKE](#)

Abstract— Solid metal/liquid metal partition coefficients for Ag and Pd were determined experimentally as a function of the S concentration of the metallic liquid. Silver is incompatible in solid metal and strongly sensitive to the S content of the metallic liquid; partition coefficients for Ag decrease more than an order of magnitude with increasi...

[Cite](#)

[Request full-text](#)

Accessory silicate mineral assemblages in the Bilanga diogenite: A petrographic study

[Article](#)

- Jan 2010
- [Kenneth Domanik](#)
- [Serena Kolar](#)
- [Donald Musselwhite](#)
- [Michael J. Drake](#)

Abstract— The petrographic relationships in diogenites between orthopyroxene and minor phases such as chromite, troilite, diopside, plagioclase, and silica are often obscured by the intense brecciation that characterizes these meteorites. Although brecciated, Bilanga preserves numerous clasts displaying primary textural relations between orthopyrox...

[Cite](#)

[Request full-text](#)

Metal/Silicate Partitioning of Mo, W and V: Effects of T, P, X and fO₂

[Article](#)

- Dec 2009
- [Eddy Hill](#)
- [K. J. Domanik](#)
- [M. J. Drake](#)

Existing studies have produced some confusion concerning the trends found for Mo and W metal/silicate partitioning. The behavior of V, however, is reasonably understood and hence allows for cross-laboratory comparisons. We are undertaking high-pressure, high-temperature experiments to study the metal/silicate partitioning behavior of all three elem...

[Cite](#)

[Request full-text](#)

Water in the Inner Solar System: Insights from Atomistic and Electronic-Structure Calculations

Article

- Sep 2009
- [K. Muralidharan](#)
- [M. Stimpfl](#)
- [Nora H. de Leeuw](#)
- [...]
- [M. J. Drake](#)

Cite

[Request full-text](#)

Silicon Mantle/Core Fractionation and the Origin of Pallasites

Article

- Sep 2009
- [Eddy Hill](#)
- [K. Domanik](#)
- [G. R. Huss](#)
- [M. J. Drake](#)

Cite

[Request full-text](#)

2009 Service Award for Derek Sears

Article

- Jul 2009
- [Michael J. Drake](#)

Cite

[Request full-text](#)

Some - Perhaps Most - Water in the Earth Must Result from Adsorption on to Grains in the Accretion Disk

Article

- Mar 2009
- [K. Muralidharan](#)
- [M. Stimpfl](#)

- [Nora H. de Leeuw](#)
- [\[...\]](#)
- [M. J. Drake](#)

We show that adsorption of water onto grains in the accretion disk must be a significant source of Earth's water. Using density functional theory we show that HDO may be preferentially retained relative to H₂O in adsorption/desorption kinetics.

[Cite](#)

[Request full-text](#)

[Experimental studies of metal–silicate partitioning of Sb: Implications for the terrestrial and lunar mantles](#)

[Article](#)

- Mar 2009
- [Kevin Righter](#)
- [Munir Humayun](#)
- [A.J. Campbell](#)
- [\[...\]](#)
- [M.J. Drake](#)

The terrestrial mantle has a well defined Sb depletion of $\sim 7 \pm 1$ (Jochum and Hofmann, 1997), and the lunar mantle is depleted relative to the Earth by a factor of $\sim 50 \pm 5$ (Wolf and Anders, 1980). Despite these well defined depletions, there are few data upon which to evaluate their origin—whether due to volatility or core formation. We have carried...

[Cite](#)

[Request full-text](#)

[Melosh Receives 2008 Harry H. Hess Medal](#)

[Article](#)

- Jan 2009
- [Michael J. Drake](#)
- [Jay Melosh](#)

H. Jay Melosh was awarded the 2008 Harry H. Hess Medal at the AGU Fall Meeting Honors Ceremony, held 17 December 2008 in San Francisco, Calif. The medal is for "outstanding achievements in research in the constitution and evolution of Earth and sister planets."

[Cite](#)

[Request full-text](#)

14. Oxygen Isotopic Composition and Chemical Correlations in Meteorites and the Terrestrial Planets

Chapter

- Dec 2008
- [David W. Mittlefehldt](#)
- [Robert N. Clayton](#)
- [Michael J. Drake](#)
- [Kevin Righter](#)

[Cite](#)

[Request full-text](#)

Origin of water in the inner Solar System: A kinetic Monte Carlo study of water adsorption on forsterite

Article

- Dec 2008
- [Krishna Muralidharan](#)
- [Pierre Deymier](#)
- [M. Stimpfl](#)
- [\[...\]](#)
- [Michael J. Drake](#)

The origin of water in the inner Solar System is not well understood. It is believed that temperatures were too high in the accretion disk in the region of the terrestrial planets for hydrous phases to be thermodynamically stable. Suggested sources of water include direct adsorption of hydrogen from the nebula into magma oceans after the terrestria...

[Cite](#)

[Request full-text](#)

Metal/olivine partition coefficients and the cooling rates of pallasites

Article

- Jul 2008
- [Eddy Hill](#)
- [K. Domanik](#)
- [M. J. Drake](#)

[Cite](#)

[Request full-text](#)

Oxygen Isotopic Composition and Chemical Correlations in Meteorites and the Terrestrial Planets

Article

- Jan 2008
- [David W. Mittlefehldt](#)
- [Robert N. Clayton](#)
- [Michael J. Drake](#)
- [Kevin Righter](#)

Recent models attempting to explain non-mass-dependent oxygen isotopic anomalies in meteorites and planets posit that they may have originated within the gas phase of the solar nebula, which suggests the potential for correlations of non-mass-dependent oxygen isotopic anomalies with other chemical fractionations generated during cooling and condens...

Cite

[Request full-text](#)

Metal/Olivine Partitioning Experiments and the Cooling Rates of Pallasites

Article

- Aug 2007
- [Eddy Hill](#)
- [M. J. Drake](#)
- [K. Domanik](#)

Cite

[Request full-text](#)

Adsorption of Water Gas on to Grains in the Accretion Disk as a Source of Inner Solar System Water: Direct Evidence from Monte Carlo Simulation

Article

- Jul 2007
- [K. Muralidharan](#)
- [P. A. Deymier](#)
- [M. Stimpfl](#)
- [\[...\]](#)
- [Nora H. de Leeuw](#)

Cite

[Request full-text](#)

Thermal Observations of OSIRIS target 1999 RQ36

Article

- Mar 2007
- [Joshua Emery](#)
- [Michael Drake](#)
- [Yanga Fernandez](#)
- [...]
- [Dante Lauretta](#)

We propose to observe the near-Earth asteroid (101955) 1999 RQ36, the target of the OSIRIS sample return mission, with IRS and IRAC. These observations will provide characterization of the composition, structure, and thermophysical properties of this distinctive asteroid. Compositionally diagnostic emissivity features measured by IRS at two longitu...

[Cite](#)

[Request full-text](#)

Water Matters

Article

- Feb 2007
- [M. J. Drake](#)
- [M. Stimpfl](#)

The origin of water on Earth and the other rocky planets is uncertain. There are problems with commonly accepted sources such as comets and asteroids. We explore adsorption of water onto grains in the accretion disk as a source.

[Cite](#)

[Request full-text](#)

Water in the Accretion Disk: Effect of Composition and Surface Structure on the Energy of Adsorption onto Olivine Grains

Article

- Feb 2007
- [M. Stimpfl](#)
- [Nora H. de Leeuw](#)
- [M. J. Drake](#)
- [P. Deymier](#)

The surface of olivine presents a variety of bonding environments for the adsorbed molecular water. Binding energies range from <44 kJ/mole to >200 kJ/mole. These interactions are strong enough to retain water on the surfaces at high temperature.

Cite

[Request full-text](#)

[Petrology and geochemistry of the NWA 3368 Eucrite](#)
[Article](#)

- Sep 2006
- [K.G. Gardner](#)
- [D. S. Lauretta](#)
- [D. H. Hill](#)
- [\[...\]](#)
- [M. J. Drake](#)

We report the petrology and geochemistry of NWA 3368, a new non-cumulate, monomict eucrite breccia with a variety of clast sizes and a pink-tinted matrix. Analytical techniques include electron microprobe, INAA, and ICP-MS.

Cite

[Request full-text](#)

[Can the Water Pressure in the Accretion Disk Sustain Water Adsorption on Olivine?](#)
[Article](#)

- Sep 2006
- [Nora H. de Leeuw](#)
- [M. Stimpfl](#)
- [Richard Richard A Catlow](#)
- [\[...\]](#)
- [A. W. Walker](#)

Cite

[Request full-text](#)

[An ångström-sized window on the origin of water in the inner solar system: Atomistic simulation of adsorption of water on olivine](#)
[Article](#)

- Aug 2006
- [M. Stimpfl](#)
- [Auriana Miranda Walker](#)
- [M.J. Drake](#)
- [\[...\]](#)

- [P. Deymier](#)

The origin of water in the inner solar system is not yet well understood. Because of the coexistence of water and small solid particulates in the accretion disk from which our planetary system formed, we propose that adsorption of water onto the surfaces/pores of forsterite could play an important role in the delivery of water to the rocky planets...

[Cite](#)

[Request full-text](#)

[Effect of composition on adsorption of water on perfect olivine surfaces](#)

[Article](#)

- Aug 2006

- [M. Stimpfl](#)
- [M.J. Drake](#)
- [Nora H. de Leeuw](#)
- [...]
- [Auriana Miranda Walker](#)

[Cite](#)

[Request full-text](#)

[The earliest history of the Earth](#)

[Article](#)

- Aug 2006

- [M. J. Drake](#)

[Cite](#)

[Request full-text](#)

[Experimental study of platinum solubility in silicate melt to 14GPa and 2273K: Implications for accretion and core formation in Earth](#)

[Article](#)

- May 2006

- [Werner Ertel-Ingrisch](#)
- [Michael J. Walter](#)
- [Michael J. Drake](#)
- [Paul Joseph Sylvester](#)

We determined the solubility limit of Pt in molten haplo-basalt (1 atm anorthite-diopside eutectic composition) in piston-cylinder and multi-anvil experiments at pressures between 0.5 and 14 GPa and temperatures from 1698 to 2223 K. Experiments were internally buffered at $\sim IW + 1$. Pt concentrations in quenched-glass samples were measured by laser-a...

[Cite](#)

[Request full-text](#)

In the Beginning There Was Water and Dust: A Look into Adsorption as a Mechanism to Explain Water in the Inner Solar System

[Article](#)

- Feb 2006
- [M. Stimpfl](#)
- [Nora H. de Leeuw](#)
- [P. Deymier](#)
- [...]
- [Auriana Miranda Walker](#)

Atomistic techniques are employed to study the interaction between water and olivine surfaces with the aim to explore if water gas adsorbed onto the dust in the accretion disk could be a possible source for water in the inner solar system.

[Cite](#)

[Request full-text](#)

Water in the Early Earth

[Article](#)

- Jan 2006
- [Yutaka Abe](#)
- [Eiji Ohtani](#)
- [Takuo Okuchi](#)
- [...]
- [M. Drake](#)

In this chapter we discuss the behavior of water in the early Earth. Earth likely formed through accretion of water-bearing planetesimals. An H₂O-rich proto-atmosphere should have formed during accretion by degassing from planetesimals and/or gravitational attraction of solar-nebula gas, and it may be a direct ancestor of Earth's present atmosphere...

[Cite](#)

[Request full-text](#)

Compositional Relationships Between Meteorites and Terrestrial Planets

Article

- Jan 2006
- [Kevin Righter](#)
- [M. J. Drake](#)
- [Edward R. D. Scott](#)

Terrestrial planets exhibit a range of core sizes, mantle, and surficial magma compositions. Determining the bulk compositions of terrestrial planets presents a challenge faced by geochemistry and cosmochemistry. Traditionally, chondritic meteorites have been called upon as suitable bulk compositions. Detailed examination of such building blocks, h...

[Cite](#)

[Request full-text](#)

[A review of meteorite evidence for the timing of magmatism and of surface or near-surface liquid water on Mars](#)

Article

Full-text available

- Dec 2005
- [Lars E. Borg](#)
- [Michael J Drake](#)

1] There is widespread photogeological evidence for ubiquitous water flowing on the surface of Mars. However, the age of surface and near-surface water cannot be deduced with high precision from photogeology. While there is clear evidence for old and young fluvial features in the photogeologic record, the uncertainty in the absolute calibration of...

[Cite](#)

[Download full-text](#)

[Melting of Allende at Pressures Between 1 and 25 Mpa](#)

Article

- Sep 2005
- [Steven Singletary](#)
- [Timothy L Grove](#)
- [M. J. Drake](#)

[Cite](#)

[Request full-text](#)

Bulk Composition of the Moon: 2. Volatiles and Isotopes

Article

- Sep 2005
- [M. J. Drake](#)
- [G. J. Taylor](#)

[Cite](#)

[Request full-text](#)

Origin of water on the terrestrial planets

Article

- Aug 2005
- [Michael J. Drake](#)
- [Humberto Campins](#)

We examine the origin of water in the terrestrial planets. We list various geochemical measurements that may be used to discriminate between different endogenous and exogenous sources of water. Late stage delivery of significant quantities of water from asteroidal and cometary sources appears to be ruled out by isotopic and molecular ratio consider...

[Cite](#)

[Request full-text](#)

The Leonard medal address: Origin of water in the terrestrial planets

Article

- Apr 2005
- [Michael J. Drake](#)

I examine the origin of water in the terrestrial planets. Late-stage delivery of water from asteroidal and cometary sources appears to be ruled out by isotopic and molecular ratio considerations, unless either comets and asteroids currently sampled spectroscopically and by meteorites are unlike those falling to Earth 4.5 Ga ago, or our measurements...

[Cite](#)

[Request full-text](#)

Influence of Silicate Melt Composition on Metal/Silicate Partitioning of W, Ge, Ga and Ni

[Article](#)

[Full-text available](#)

- Feb 2005
- [Steven Singletary](#)
- [K. Domanik](#)
- [M. J. Drake](#)

The depletion of the siderophile elements in the Earth's upper mantle relative to the chondritic meteorites is a geochemical imprint of core segregation. Therefore, metal/silicate partition coefficients ($D_{m/s}$) for siderophile elements are essential to investigations of core formation when used in conjunction with the pattern of elemental abundances...

[Cite](#)

[Download full-text](#)

NWA 2736: An Unusual New Graphite-bearing Aubrite

[Article](#)

- Feb 2005
- [J. J. Lowe](#)
- [D. H. Hill](#)
- [K. J. Domanik](#)
- [...]
- [Marvin Killgore](#)

Aubrites are enstatite-rich achondrites that are probably related to enstatite chondrite parent bodies. The exact link, if any, is not understood. They may have formed from

melting within one or more enstatite chondrite parent bodies. Description of NWA 2736: One complete stone weighing 171.51 grams was recovered from the Sahara Desert by nomads so...

[Cite](#)

[Request full-text](#)

Olivine and Ca-Phosphate in the Diogenites Manegaon and Roda

[Article](#)

- Feb 2005
- [K. J. Domanik](#)
- [L. C. Sideras](#)
- [M. J. Drake](#)

The textural relationships between the different primary minerals in igneous rocks provide one of the most fundamental pieces of evidence available for inferring the crystallization history of their parent magmas. Unfortunately, the high degree of brecciation that characterizes most diogenites, along with the low modal abundance and small grain siz...

[Cite](#)

[Request full-text](#)

Early Accretion of Water: Implications for the Oxidation State of the Planets

[Article](#)

- Dec 2004
- [M. Stimpfl](#)
- [M. J. Drake](#)
- [D. S. Lauretta](#)

Introduction: There are two main scenarios that account for the accretion of water to the terrestrial planets: either the water originated outside of the inner solar system and was later delivered to the terrestrial planets by means of some suitable mechanism, or the source of water was local, i.e. it came from the same feeding zone of the Earth's...

[Cite](#)

[Request full-text](#)

Metal/Silicate Partitioning of W, Ge, Ga and Ni: Dependence on Silicate Melt Composition

[Article](#)

- Dec 2004

- [Steven Singletary](#)
- [M. J. Drake](#)

Metal/silicate partition coefficients ($D_{m/s}$) for siderophile elements are essential to investigations of core formation when used in conjunction with the pattern of elemental abundances in the Earth's mantle (Drake and Righter, 2002; Jones and Drake, 1986; Righter et al. 1997). The partitioning of siderophile elements is controlled by temperature,...

[Cite](#)

[Request full-text](#)

Origin of water in the terrestrial planets

[Article](#)

- Aug 2004
- [M.J. Drake](#)

I examine the origin of water in the terrestrial planets. Late-stage delivery of water from asteroidal and cometary, sources appears to be ruled out by isotopic and molecular ratio considerations, unless either comets and asteroids currently sampled spectroscopically and by meteorites are unlike those falling to Earth 4.5 Ga ago, or our measurement...

[Cite](#)

[Request full-text](#)

Adsorption as a Mechanism to Deliver Water to the Earth

[Article](#)

- Aug 2004
- [M. Stimpfl](#)
- [D. S. Laurretta](#)
- [M. J. Drake](#)

[Cite](#)

[Request full-text](#)

Metal-Silicate Partitioning of Volatile, Siderophile Elements: New Results for Sb and As

[Article](#)

- Aug 2004
- [Kevin Righter](#)
- [A. J. Campbell](#)

- [Munir Humayun](#)
- [M. J. Drake](#)

[Cite](#)

[Request full-text](#)

Implications of Chondrule Formation in a Gas of Solar Composition [Article](#)

- Jan 2004
- [D. S. Lauretta](#)
- [M. J. Drake](#)
- [M. Stimpfl](#)

[Cite](#)

[Request full-text](#)

How did the Terrestrial Planets Acquire Their Water? [Article](#)

- Jan 2004
- [M. J. Drake](#)
- [M. Stimpfl](#)
- [D. S. Lauretta](#)

[Cite](#)

[Request full-text](#)

Partition Coefficients at High Pressure and Temperature [Article](#)

- Nov 2003
- [Kevin Righter](#)
- [M. J. Drake](#)

Differentiation of terrestrial planets includes separation of a metallic core and possible later fractionation of mineral phases within either a solid or molten mantle (Figure 1). Lithophile and siderophile elements can be used to understand these two different physical processes, and ascertain whether they operated in the early Earth. The distribu...

[Cite](#)

[Request full-text](#)

Vanadium, Cr, and Si and the Mg/Si Ratio in the Earth

Article

- Jul 2003
- [M. J. Drake](#)
- [K. Domanik](#)
- [Kevin Righter](#)

Cite

[Request full-text](#)

Citation for presentation of the Distinguished Service Award to Denis M. Shaw

Article

- Jul 2003
- [Michael J. Drake](#)

Cite

[Request full-text](#)

Trace Elements in High-Ca Pyroxene and Plagioclase in the Bilanga Diogenite: Implications for the Magmatic Evolution of Diogenites

Article

- Feb 2003
- [K. J. Domanik](#)
- [C. K. Shearer](#)
- [Justin Hagerty](#)
- [...]
- [M. J. Drake](#)

High-Ca pyroxene and plagioclase are typically present as minor phases in diogenites. However, although the trace element content of diogenite orthopyroxene has been measured in a number of studies; almost no trace element data is available for the high-Ca pyroxene and plagioclase with which it routinely coexists in these meteorites. These data cou...

Cite

[Request full-text](#)

Bilanga: A Unique Diogenite

Article

- May 2002

- [S. E. Kolar](#)
- [K. J. Domanik](#)
- [D. S. Musslewhite](#)
- [M. J. Drake](#)

The Bilanga diogenite is petrographically examined and found to contain diopside and plagioclase aggregates up to several 100 microns in size. Small scale, original igneous contacts between these phase are often preserved despite brecciation. Additional information is contained in the original extended abstract.

[Cite](#)

[Request full-text](#)

[Petrology of Antarctic Eucrites PCA 91078 and PCA 91245](#)

[Article](#)

- May 2002
- [L. M. Howard](#)
- [K. J. Domanik](#)
- [M. J. Drake](#)
- [David W. Mittlefehldt](#)

Antarctic eucrites PCA 91078 and PCA 91245, are petrographically characterized and found to be unpaired, type 6, basaltic eucrites. Observed textures that provide insight into the petrogenesis of these meteorites are also discussed. Additional information is contained in the original extended abstract.

[Cite](#)

[Request full-text](#)

[What is the Earth Made Of?](#)

[Article](#)

[Full-text available](#)

- May 2002
- [M. J. Drake](#)
- [Kevin Righter](#)

Measurements made on materials from Earth, Mars, comets, and meteorites of Mg/Si, Al/Si, oxygen, Os, D/H, Ar/H₂O, and Kr/Xe ratios, all lead to the conclusion that no primitive material similar to Earth mantle material is currently represented in our meteorite collections. Additional information is contained in the original extended abstract.

[Cite](#)

[Download full-text](#)

Determining the composition of the Earth

Article

Full-text available

- Apr 2002
- [Michael J Drake](#)
- [Kevin Righter](#)

A long-standing question in the planetary sciences asks what the Earth is made of. For historical reasons, volatile-depleted primitive materials similar to current chondritic meteorites were long considered to provide the 'building blocks' of the terrestrial planets. But material from the Earth, Mars, comets and various meteorites have Mg/Si and Al...

[Cite](#)

[Download full-text](#)

Narrow horizons in astrobiology

Article

- Mar 2002
- [Michael J Drake](#)
- [Bruce M Jakosky](#)

Mission planners must realize that astrobiology is more than a search for life

[Cite](#)

[Request full-text](#)

Experimental Studies of Metal-Silicate Partitioning of Sb

Article

- Sep 2001
- [Kevin Righter](#)
- [D. Hill](#)
- [J. Collins](#)
- [...]

- [M. J. Drake](#)

[Cite](#)

[Request full-text](#)

[Constraints on the Depth of an Early Terrestrial Magma Ocean](#)

[Article](#)

[Full-text available](#)

- [Aug 2001](#)
- [Kevin Righter](#)
- [M. J. Drake](#)

[Cite](#)

[Download full-text](#)

[Presidential Address: Presented 2000 August 28, Chicago, Illinois, USA The eucreite/Vesta story](#)

[Article](#)

- [Apr 2001](#)
- [Michael J. DRAKE](#)

Abstract— Many lines of evidence indicate that meteorites are derived from the asteroid belt but, in general, identifying any meteorite class with a particular asteroid has been problematical. One exception is asteroid 4 Vesta, where a strong case can be made that it is the ultimate source of the howardite-eucrite-diogenite (HED) family of basaltic...

[Cite](#)

[Request full-text](#)

[Experimentally Determined Solubilities of Pt up to 90 Kbars and 1850°C](#)

[Article](#)

- [Jan 2001](#)
- [Werner Ertel-Ingrisch](#)
- [M. J. Walter](#)
- [Paul Joseph Sylvester](#)
- [M. J. Drake](#)

PGEs are indicators of planetary differentiation processes based on their siderophilicity. Results of our HP experiments (5 to 90 kbar; 1400 to 1850°C) indicate that P does not influence the Pt partitioning, supporting the late veneer hypothesis.

Cite

[Request full-text](#)

Metal-silicate partitioning of Co, Ga, and W: Dependence on silicate melt composition

[Article](#)

- Nov 2000
- [Lv Windy](#)
- [Michael J. Drake](#)

This study investigates the effect of silicate melt composition on metal/silicate partitioning for Co^{2+} , Ga^{3+} , and W^{4+} at 1300°C , 1 atm, and a $\log f_{\text{O}_2}$ of -12 . Five glasses in the system $\text{MgO}-\text{CaO}-\text{Al}_2\text{O}_3-\text{TiO}_2-\text{SiO}_2$ with nbo/t (nonbridging oxygens/tetrahedrally coordinated cations) values ranging from 0.25 to 1.52 were used as starting materials. For W a...

Cite

[Request full-text](#)

Early Outgassing of Mars: Implications from Experimentally Determined Solubility of Iodine in Silicate Magmas

[Article](#)

- Nov 2000
- [Donald S. Musselwhite](#)
- [Michael J. Drake](#)

We have determined experimentally the solubility of iodine in a range of synthetic basaltic (CMAS) liquids. The solubility of iodine ranges from 1.2×10^{-4} to 2.6×10^{-3} g/g-atm (0.02 to 0.45 ccSTP/g-atm) over the range of compositions studied. These values are a few orders of magnitude greater than the solubility of xenon in silicate liquids. This dif...

Cite

[Request full-text](#)

Metal/silicate equilibrium in the early Earth-New constraints from the volatile moderately siderophile elements Ga, Cu, P, and Sn

[Article](#)

- Oct 2000
- [Kevin Righter](#)
- [Michael J Drake](#)

We report new measurements of metal/silicate partition coefficients for Ga, Cu, and Sn at 10 to 90 kb and 1250 to 1900°C. We show that all three of these siderophile elements ($D(\text{metal/silicate}) > 10$) become more lithophile at high pressure and temperature conditions. Metal/silicate partition coefficients calculated for the conditions of an early de...

[Cite](#)

[Request full-text](#)

Accretion and primary differentiation of the Earth: A personal journey

[Article](#)

- Jul 2000
- [Michael J Drake](#)

The accretion of the Earth was a violent series of events dominated by the addition of objects one third to one tenth of the mass of the growing planet. During the later stages of accretion, these collisions deposited enough energy to at least partly melt the Earth, possibly multiple times. The result was an ocean of magma. Metal sank through this...

[Cite](#)

[Request full-text](#)

Crystallization of magmatic iron meteorites: The effects of phosphorus and liquid immiscibility

[Article](#)

- Jul 2000
- [Nancy Chabot](#)
- [M.J. Drake](#)

Magmatic iron meteorites are commonly thought to have formed by fractional crystallization of the metallic cores of asteroid-sized bodies. As fractional crystallization proceeds, light elements such as P and S become enriched in the molten portion of the core. The light element content of the metallic liquid influences the partitioning behavior of...

[Cite](#)

[Request full-text](#)

Accretion of the Earth: Implications from Experimental Constraints: Pt

[Article](#)

- Mar 2000
- [Werner Ertel-Ingrisch](#)
- [Paul Joseph Sylvester](#)

- [M. J. Drake](#)

[Cite](#)

[Request full-text](#)

Potassium solubility in metal: The effects of composition at 15 kbar and 1900°C on partitioning between iron alloys and silicate melts

[Article](#)

- Oct 1999
- [Nancy Chabot](#)
- [Michael J Drake](#)

To determine the role of radioactive heating as an energy source in planetary cores, the solubility of K in metal has been examined experimentally. All experiments were conducted at 15 kbar and 1900°C and involved K partitioning between Fe alloys and silicate melts. Experiments conducted with different concentrations of S in the metallic liquid ind...

[Cite](#)

[Request full-text](#)

Siderophile geochemistry of Ga, Ge, and Sn: Cationic oxidation states in silicate melts and the effect of composition in iron-nickel alloys

[Article](#)

- Oct 1999
- [Christopher J. Capobianco](#)
- [Michael J. Drake](#)
- [Jessie de'Aro](#)

We report a series of metal–silicate partitioning experiments for Ga, Ge, and Sn to characterize the dependence of the partition coefficient, D, on oxygen fugacity, fO_2 . These were isothermal (1260°C) and isobaric (1 bar) experiments using a silicate composition that approximates a eucritic meteorite. It is well known that elements such as Ni, whic...

[Cite](#)

[Request full-text](#)

Effect of water on metal-silicate partitioning of siderophile elements: A high pressure and temperature terrestrial magma ocean and core formation

[Article](#)

- Sep 1999

- [Kevin Righter](#)
- [Michael J. Drake](#)

Recent proposals of metal-silicate equilibrium at the base of a deep hydrous magma ocean are based on experimental data obtained under anhydrous conditions. We have undertaken a series of experiments at 10 kbar and 1300°C, designed to isolate the effect of dissolved water on the partitioning of the siderophile elements Ni, Co, Mo, W, and P between...

[Cite](#)

[Request full-text](#)

Crystallization of Magmatic Iron Meteorites and Liquid Immiscibility in the Iron-Phosphorus-Sulfur System

[Article](#)

- Jun 1999
- [Nancy Chabot](#)
- [M. J. Drake](#)

[Cite](#)

[Request full-text](#)

Scientists debate origin of Earth and Moon

[Article](#)

- May 1999
- [Alex N. Halliday](#)
- [Michael J. Drake](#)

The big questions of how the Earth and Moon formed and first developed have, for many years, been at the borderline of tractability because of what appears to be an almost complete obliteration of any physical, chemical, or isotopic record of the first 500 million years of Earth history. We do know that in the first 200 million years the Earth acqu...

[Cite](#)

[Request full-text](#)

Colliding Theories

[Article](#)

- Mar 1999
- [Alex N. Halliday](#)
- [Michael J. Drake](#)

When researchers try to find out how the Earth and the moon formed, they quickly come up against the problem that direct evidence is impossible to come by. Halliday and Drake discuss recent progress towards understanding Earth and moon formation using simulations, high pressure experiments and isotope geochemistry. The most widely accepted theory a...

[Cite](#)

[Request full-text](#)

Partitioning of W Between Liquid Metal, Solid Silicates, and Liquid Silicates at High Pressures and Temperatures: Implications for the ^{182}W Isotope Anomalies in Lunar and Martian Samples

[Article](#)

- Mar 1999
- [Kevin Righter](#)
- [M. J. Drake](#)

New high pressure and temperature experiments on partially molten basalt and chondrite show that the ^{182}W anomalies measured in lunar and martian samples must be interpreted with a full understanding of metal-silicate and clinopyroxene-silicate liquid partitioning.

[Cite](#)

[Request full-text](#)

Experimental Constraints on the Accretion of the Earth: Pt

[Article](#)

- Mar 1999
- [Werner Ertel-Ingrisch](#)
- [Paul Joseph Sylvester](#)
- [M. J. Drake](#)

Pt equilibrium solubilities in An-Di eutectic silicate melt was determined at 10 to 20 kbars and 1450o to 1850oC in piston cylinder experiments. Results are compared to previous work at 1 atm and 2 to 4 kbar.

[Cite](#)

[Request full-text](#)

Crystallization of magmatic iron meteorites: The role of mixing in the molten core

[Article](#)

- Mar 1999
- [Nancy Chabot](#)

- [M.J. Drake](#)

The IIIAB group is the largest of the magmatic iron meteorite groups and consequently is commonly used to test models of asteroid core crystallization. Simple fractional crystallization calculations appear to reproduce the general shape of the elemental trends observed in the IIIAB group when these trends are plotted vs. Ni, as is traditionally done...

[Cite](#)

[Request full-text](#)

Potassium solubility in metal: the effects of composition at 15kbar and 1900 on partitioning between iron alloys and silicate melts

[Article](#)

- Jan 1999
- [Nancy Chabot](#)
- [M.J. Drake](#)

[Cite](#)

[Request full-text](#)

The Effect of Silicate Composition on the Solubility of K in Metal

[Article](#)

- Jan 1999
- [Nancy Chabot](#)
- [M. J. Drake](#)

[D(K)] metal/silicate increases by ~100 with increasing depolymerization of the silicate melt. At 15 kb and 1900°C D(K) for the Earth yields a present day heat generation ~100 lower than needed to drive the Earth's geodynamo. Higher P and T will be studied.

[Cite](#)

[Request full-text](#)

Examining Crystallization Scenarios for Magmatic Iron Meteorites with a Mixing Model

[Article](#)

- Jul 1998
- [Nancy Chabot](#)
- [M. J. Drake](#)

Cite

[Request full-text](#)

Uranium in the Lunar Crust: Implications for Lunar Origin and Evolution
[Article](#)

- Jul 1998
- [M. J. Drake](#)
- [J. P. Lowe](#)
- [Kevin Righter](#)
- [\[...\]](#)
- [P. E. Clark](#)

Cite

[Request full-text](#)

Ion Microprobe Study of Plagioclase-Basalt Partition Experiments at Natural Concentration Levels of Trace Elements
[Article](#)

- Apr 1998
- [Ilya Bindeman](#)
- [Andrew M. Davis](#)
- [Michael J. Drake](#)

We present here a study of plagioclase/melt partitioning of trace elements at their natural concentration levels, using sample charges from the widely cited plagioclase/melt partitioning experiments of Drake and Weill (1975). In these experiments, sample charges were doped to ~1 wt% with Sr, Ba, rare earth elements (REE) and Y, but each charge was...

Cite

[Request full-text](#)

Metal-Silicate Equilibrium in the Early Earth -- New Constraints from Volatile Moderately Siderophile Elements Ga, Sn and Cu
[Article](#)

- Mar 1998
- [Kevin Righter](#)
- [M. J. Drake](#)

Cite

[Request full-text](#)

The Effect of S on the Solubility of K in Metal
[Article](#)

- Mar 1998
- [Nancy Chabot](#)
- [M. J. Drake](#)

[Cite](#)

[Request full-text](#)

Determination of RE Contents in Experimental Glasses by Two Techniques: Ion Microprobe and Neutron Activation Analysis
[Article](#)

- Jan 1998
- [Kevin Righter](#)
- [D. J. Lindstrom](#)
- [R. R. Martinez](#)
- [...]
- [M. J. Drake](#)

[Cite](#)

[Request full-text](#)

An experimental study of silver and palladium partitioning between solid and liquid metal, with applications to iron meteorites
[Article](#)

- Sep 1997
- [Nancy Chabot](#)
- [M.J. Drake](#)

Solid metal/liquid metal partition coefficients for Ag and Pd were determined experimentally as a function of the S concentration of the metallic liquid. Silver is incompatible in solid metal and strongly sensitive to the S content of the metallic liquid; partition coefficients for Ag decrease more than an order of magnitude with increasing S conte...

[Cite](#)

[Request full-text](#)

An Experimental Investigation of Liquid Immiscibility During Crystallization of Magmatic Iron Meteorites

[Article](#)

- Jul 1997
- [Nancy Chabot](#)
- [M. J. Drake](#)

[Cite](#)

[Request full-text](#)

Prediction of siderophile element metal-silicate partition coefficients to 20 GPa and 2800°C: The effects of pressure, temperature, oxygen fugacity, and silicate and metallic melt compositions

[Article](#)

[Full-text available](#)

- Mar 1997
- [Kevin Righter](#)
- [M.J. Drake](#)
- [Gregory M Yaxley](#)

We report new metal-silicate partition coefficients for Ni, Co and P at 7.0 GPa (1650–1750°C), and Ni, Co, Mo, W and P at 0.8, 1.0 and 1.5 GPa (1300–1400°C). Guided by thermodynamics, all available metal-silicate partition coefficients, $D(i)$, where i is Ni, Co, P, Mo and W, are regressed against $\ln(fO_2)$, $\ln(1 - X_s)$ (X_s is mole fraction of S in...

[Cite](#)

[Download full-text](#)

Formation of eucrites and diogenites on a Vesta-sized asteroid. I - Core formation

[Article](#)

- Feb 1997
- [Kevin Righter](#)
- [Michael J. Drake](#)

Core formation in asteroid-sized planetesimals occurred early (less than 15 Ma after T(0)) as has been recently demonstrated using W-182/W-184 ratios. Depletion of siderophile elements in the mantle of the HED parent body has been cited as evidence that metal had segregated into a core before the eucrites formed. Calculations and experiments indica...

Cite

[Request full-text](#)

1

2

3

Citations

... The early Earth experienced at least one episode of extensive silicate melting, also known as magma ocean (e.g., Abe and Matsui, 1988; Abe, 1993; Solomatov and Stevenson, 1993a; Abe, 1997; Solomatov, 2000; Drake, 2000; Elkins-Tanton, 2012). upwards until it reaches the surface of the planet, and the BMO front progresses downwards until it reaches the CMB. ...

Reference: [Timescales of chemical equilibrium between the convecting solid mantle and over-/underlying magma oceans](#)

[Accretion and primary differentiation of the Earth: A personal journey](#)

[Citing article](#)

- Jul 2000
- [M. J. Drake](#)

[View](#)

... Primitive NEAs have and continue to be targeted by spacecraft, including robotic and human space missions (e.g., Abell et al. 2015, Mazanek et al., 2016. The two primitive NEAs that are about to be visited by sample-return missions are (101955) Bennu, target of NASA's OSIRIS-REx (Lauretta et al., 2010), and (162173) Ryugu, target of JAXA's Hayabusa2 (Tsuda et al., 2014). Both missions will rendezvous their respective asteroids in 2018. ...

Reference: [Compositional Diversity Among Primitive Asteroids](#)

[Asteroid \(101955\) 1999 RQ36: Optimum Target for an Asteroid Sample Return Mission](#)

[Citing article](#)

- Jul 2010
- [D. S. Lauretta](#)
- [M. J. Drake](#)
- [R. P. Benz](#)
- [H. Campins](#)

- [Daniel J. Scheeres](#)

[View](#)

... Volatiles can be present in planets from the early stages of planet formation, when there is still gas in the disc; or reach the protoplanets after the gas is gone. In the former case, in principle only planets located beyond the iceline can have volatiles, although water could also be produced within a dry protoplanet from a reaction between H₂ and FeO (Ikoma and Genda, 2006), or retained in olivine grains via adsorption (Drake, 2005). ...

Reference: [Setting the Stage: Planet formation and Volatile Delivery](#)

[Origin of water in the terrestrial planets](#)

[Citing article](#)

- Aug 2004

- [M. J. Drake](#)

[View](#)

... Jones and Drake, 1983;Haack and Scott. 1993;Ulf-Møller, 1998;Chabot and Drake, 1999) may have been wrongly assuming that effects from subsolidus schreibersite evolution were negligible on the Ni solid metal composition. While Ni is clearly affected by schreibersite growth, whether the effects on other elements are indeed negligible or not is less clear. ...

Reference: [Experimental Partitioning of Trace Elements into Schreibersite with Applications to IIG Iron Meteorites](#)

[Crystallization of magmatic iron meteorites: The role of mixing in the molten core](#)

[Citing article](#)

- Mar 1999

- [N. L. Chabot](#)
- [M. J. Drake](#)

[View](#)

... Morbidelli et al. (2000) proposed that the bulk of Earth's water was accreted from a few water-rich embryos from the outer asteroid region, beyond 2-2.5 AU (a lesser amount of water, according to Morbidelli et al. (2000), was accreted as well from smaller bodies in the asteroid belt and comets). The water content of protoplanets 1 in the habitable zone is assumed to be very small, based on a number of considerations (Lunine, 2006; but see Drake and Righter, 2002), so an external source of water is needed. The work of Morbidelli et al. (2000), as well as earlier authors (e.g., Wetherill, 1996), showed that planetary accretion is a stochastic process, especially if much of the mass is contained in a small number of large embryos. ...

Reference: [High-Resolution Simulations of The Final Assembly of Earth-Like Planets. 2. Water Delivery And Planetary Habitability](#)

What is the Earth Made Of?

Citing article

- May 2002
- NTRS
- M. J. Drake
- K. Righter

View full-text

... Papoular suggests that in diffuse clouds, the grain temperature is too high for water to stick to carbonaceous grains, for which calculations predict adsorption energies of 2–9 kJ mol⁻¹ on the bare surface (Papoular, 2005). However, water has a much higher interaction energy with silicate grain surfaces, (Stimpfl et al., 2006) where, in principle, ice formation could be easily nucleated. That water ice is not observed is thus probably due to the interstellar radiation field which penetrates further into diffuse clouds, leading to photodissociation of any adsorbed water (Cuppen and Herbst, 2007; Williams et al., 1992). ...

Reference: [How Hydroxylation Affects Hydrogen Adsorption and Formation on Nanosilicates](#)

Effect of composition on adsorption of water on perfect olivine surfaces

Citing article

- Aug 2006
- M. Stimpfl
- M. J. Drake
- Nora H. de Leeuw
- P. A. Deymier
- AM Walker

View

... Diogenites consist mainly of orthopyroxene with or without olivine. They are usually coarse-grained and have Mg/(Mg + Fe) values higher than eucrites, suggesting that they were derived from the lower crust or the mantle of Vesta, based on magma ocean models [3][4][5][6][7][8]. Howardites are mechanical mixtures of eucrite and diogenite components, representing materials from the surface of Vesta.

...

Reference: [Evidence of metasomatism in the interior of Vesta](#)

A magma ocean on Vesta: Core formation and petrogenesis of eucrites and diogenites

Citing article

- Feb 2010

- [K. Righter](#)
- [M. J. Drake](#)

[View](#)

... Spitzer thermal emission data provide firm constraints on the average regolith grain size (Fig. 7a; Emery et al. 2014). Regolith grains that are comparable in size to the thermal skin depth would behave like bedrock. ...

Reference: [The OSIRIS-REx target asteroid \(101955\) Bennu: Constraints on its physical, geological, and dynamical nature from astronomical observations](#)

Thermal and Physical Characterization of the OSIRIS-REx Target Asteroid (101955) 1999 RQ36

[Citing article](#)

- Oct 2012

- [J. P. Emery](#)
- [M. S. Kelley](#)
- [Y. R. Fernandez](#)

- [C. W. Hergenrother](#)
- [M. J. Drake](#)

[View](#)

... On the ongoing debate, terrestrial planets are argued to have accreted from different classes of chondrites (i.e. carbonaceous (CC), ordinary (OC), and enstatite chondrites (EC)), achondrites or no longer extant meteorites combining in variable proportions [1][2][3][4][5] . It is generally assumed that all planetary bodies were formed from primitive solar nebula that had carbonaceous chondritic composition because of the similar relative elemental abundances of CI chondrite as solar photosphere for all except the most volatile elements 6

Reference: [Si-Mg isotopes in enstatite chondrites and accretion of reduced planetary bodies](#)

Determining the composition of the Earth

[Citing article](#)

- Apr 2002
- PubMed
- [M. J. Drake](#)
- [K. Righter](#)

[View full-text](#)

... [elsevier.com/locate/pepi](#) and high temperature (e.g., Ito et al., 1993;Ohtani and Yurimoto, 1996;Chabot and Drake, 1999;Gessmann and Wood, 2002;Murthy et al., 2003;Hirao et al., 2006;Bouhifd et al., 2007;Corgne et al., 2007).

These studies reported that the partition coefficient of potassium, D_K (=the content of potassium in metal [wt%]/ the content of potassium in silicate [wt%]), could vary from 0.0047 to 2.4, depending on the effects of temperature, pressure, oxygen fugacity, and metal and silicate compositions (e.g., NBO/T; the number of non-bridging oxygen divided by the number of tetrahedral cations). ...

Reference: [The abundance of potassium in the Earth's core](#)

Potassium solubility in metal: the effects of composition at 15kbar and 1900 on partitioning between iron alloys and silicate melts

[Citing article](#)

- Jan 1999
- [N. L. Chabot](#)
- [M. J. Drake](#)

[View](#)

Top co-authors (50)

[Kevin Righter](#)

- NASA

[M. Stimpfl](#)

- The University of Arizona

[James H Jones](#)

- Indiana University-Purdue University Indianapolis

[Nancy Chabot](#)

- Johns Hopkins University

[Allan Treiman](#)

- Universities Space Research Association

[Eilis Mcfarlane](#)

- University College Dublin

Christopher J. Capobianco

Kenneth J. Domanik

P. A. Deymier

D. S. Lauretta

Top journals

Geochimica et Cosmochimica Acta (46)

Meteoritics (30)

Meteoritics & planetary science (15)

Earth and Planetary Science Letters (9)

Affiliations

The University of Arizona
Department

- Department of Planetary Sciences
- Department of Materials Sciences and Engineering

Lunar and Planetary Institute

Universitat Oberta de Catalunya

Disciplines

Physics

Topic

- Materials Science
- Condensed Matter Physics
- Materials Physics
- Solid State Physics

Geoscience

Topic

- [Geology](#)

Chemistry

Topic

- [Physical Chemistry](#)
- [Materials Chemistry](#)
- [Analytical Chemistry](#)

Publication Stats

Citations
7,955

[Browse more researchers](#)

Join ResearchGate to find the people and research you need to help your work.

- **17+ million** members
- **135+ million** publications
- **700k+** research projects

[Join for free](#)

- [Recruit researchers](#)
 - [Join for free](#)
 - [Login](#)

Company

[About us](#)

[News](#)

[Careers](#)

Support

[Help Center](#)

Business solutions

[Advertising](#)

[Recruiting](#)

© 2008-2020 ResearchGate GmbH. All rights reserved.